

Pollinator Plot Southwest 38-541

Updated: 2021

This mix has been designed to support specialist bees, many Lepidoptera species and includes a wide range of plant families to maximize insect use, bloom periods and the long-term resiliency of the mix.



Common Name	Scientific Name	Rate (lb/ac)	% of Mix (by weight)	% by Seed	Seeds/sq ft
big bluestem	<i>Andropogon gerardii</i>	0.11	0.59%	0.88%	0.40
side-oats grama	<i>Bouteloua curtipendula</i>	1.36	7.35%	6.59%	3.00
blue grama	<i>Bouteloua gracilis</i>	0.07	0.37%	2.20%	1.00
kalm's brome	<i>Bromus kalmii</i>	0.34	1.84%	2.20%	1.00
little bluestem	<i>Schizachyrium scoparium</i>	0.54	2.94%	6.59%	3.00
Indian grass	<i>Sorghastrum nutans</i>	0.07	0.37%	0.66%	0.30
prairie dropseed	<i>Sporobolus heterolepis</i>	0.05	0.28%	0.66%	0.30
nodding wild rye	<i>Elymus canadensis</i>	0.26	1.41%	1.10%	0.50
junegrass	<i>Koeleria macrantha</i>	0.00	0.00%	0.00%	0.00
	Grasses Subtotal	2.80	15.15%	20.88%	9.50
short sedge	<i>Carex brevior</i>	0.02	0.10%	0.44%	0.20
Bicknell's sedge	<i>Carex bicknellii</i>	0.05	0.26%	0.66%	0.30
	Sedges & Rushes Subtotal	0.07	0.36%	1.10%	0.50
lead plant	<i>Amorpha canescens</i>	0.07	0.36%	0.66%	0.30
Canada milk vetch	<i>Astragalus canadensis</i>	0.03	0.17%	0.44%	0.20
partridge pea	<i>Chamaecrista fasciculata</i>	0.15	0.82%	0.33%	0.15
white prairie clover	<i>Dalea candida</i>	0.07	0.39%	1.10%	0.50
purple prairie clover	<i>Dalea purpurea</i>	0.36	1.96%	4.39%	2.00
Canada tick trefoil	<i>Desmodium canadense</i>	0.01	0.08%	0.07%	0.03
wild licorice	<i>Glycyrrhiza lepidota</i>	0.02	0.08%	0.05%	0.02
	Legumes Subtotal	0.71	3.86%	7.04%	3.20
common yarrow	<i>Achillea millefolium</i>	0.01	0.03%	0.90%	0.41
blue giant hyssop	<i>Agastache foeniculum</i>	0.03	0.17%	2.27%	1.03

Prairie Wild Onion	<i>Allium stellatum</i>	0.03	0.14%	0.22%	0.10
lead plant	<i>Amorpha canescens</i>	0.00	0.00%	0.00%	0.00
Canada anemone	<i>Anemone canadensis</i>	0.02	0.10%	0.12%	0.06
long-headed thimbleweed	<i>Anemone cylindrica</i>	0.01	0.06%	0.22%	0.10
common milkweed	<i>Asclepias syriaca</i>	0.14	0.74%	0.44%	0.20
butterfly milkweed	<i>Asclepias tuberosa</i>	0.04	0.24%	0.15%	0.07
Flodman's Thistle	<i>Cirsium flodmanii</i>	0.01	0.05%	0.05%	0.02
bird's foot coreopsis	<i>Coreopsis palmata</i>	0.02	0.12%	0.18%	0.08
tall cinquefoil	<i>Drymocallis arguta</i>	0.01	0.06%	2.20%	1.00
narrow-leaved purple coneflower	<i>Echinacea angustifolia</i>	0.06	0.32%	0.33%	0.15
bottle gentian	<i>Gentiana andrewsii</i>	0.01	0.05%	2.20%	1.00
autumn sneezeweed	<i>Helenium autumnale</i>	0.05	0.25%	4.92%	2.24
Maximilian's sunflower	<i>Helianthus maximilianii</i>	0.06	0.34%	0.66%	0.30
stiff sunflower	<i>Helianthus pauciflorus</i>	0.03	0.15%	0.09%	0.04
Early Sunflower	<i>Heliopsis helianthoides</i>	0.17	0.93%	0.88%	0.40
alumroot	<i>Heuchera richardsonii</i>	0.01	0.03%	3.53%	1.61
rough blazing star	<i>Liatris aspera</i>	0.01	0.05%	0.11%	0.05
northern plains blazing star	<i>Liatris ligulistylis</i>	0.01	0.07%	0.11%	0.05
great blazing star	<i>Liatris pycnostachya</i>	0.01	0.07%	0.11%	0.05
Winged Loosestrife	<i>Lythrum alatum</i>	0.01	0.04%	1.76%	0.80
wild bergamot	<i>Monarda fistulosa</i>	0.05	0.25%	2.65%	1.21
common evening primrose	<i>Oenothera biennis</i>	0.02	0.09%	1.25%	0.57
stiff goldenrod	<i>Oligoneuron rigidum</i>	0.06	0.34%	2.07%	0.94
wood betony	<i>Pedicularis canadensis</i>	0.01	0.03%	0.18%	0.08
large-flowered beard tongue	<i>Penstemon grandiflorus</i>	0.08	0.42%	0.88%	0.40
prairie phlox	<i>Phlox pilosa</i>	0.01	0.04%	0.11%	0.05
Virginia mountain mint	<i>Pycnanthemum virginianum</i>	0.01	0.07%	2.20%	1.00
gray-headed coneflower	<i>Ratibida pinnata</i>	0.07	0.39%	1.76%	0.80
prairie rose	<i>Rosa arkansana</i>	0.00	0.00%	0.00%	0.00
black-eyed susan	<i>Rudbeckia hirta</i>	0.07	0.37%	5.05%	2.30
Lance-leaved Figwort	<i>Scrophularia lanceolata</i>	0.02	0.08%	2.32%	1.06
field blue-eyed grass	<i>Sisyrinchium campestre</i>	0.00	0.02%	0.11%	0.05
gray goldenrod	<i>Solidago nemoralis</i>	0.01	0.05%	2.20%	1.00
showy goldenrod	<i>Solidago speciosa</i>	0.03	0.18%	2.20%	1.00
heath aster	<i>Symphotrichum ericoides</i>	0.01	0.06%	1.76%	0.80
smooth aster	<i>Symphotrichum laeve</i>	0.05	0.27%	2.20%	1.00
New England aster	<i>Symphotrichum novae-angliae</i>	0.04	0.20%	1.98%	0.90
tall meadow-rue	<i>Thalictrum dasycarpum</i>	0.07	0.40%	0.44%	0.20
bracted spiderwort	<i>Tradescantia bracteata</i>	0.03	0.18%	0.28%	0.13
hoary vervain	<i>Verbena stricta</i>	0.02	0.12%	0.50%	0.23
Culver's root	<i>Veronicastrum virginicum</i>	0.01	0.05%	6.26%	2.85
bearded birdfoot violet	<i>Viola pedatifida</i>	0.00	0.03%	0.11%	0.05
heart-leaved alexanders	<i>Zizia aptera</i>	0.04	0.20%	0.35%	0.16
golden alexanders	<i>Zizia aurea</i>	0.07	0.37%	0.62%	0.28
	Forbs Subtotal	1.53	8.22%	58.90%	26.80
Oats	<i>Avena sativa</i>	13.47	880.53%	22.38%	6.00

	Cover Crop Subtotal	13.47	880.53%	22.38%	6.00
	Total	18.51	907.76%	109.20%	45.51

Seed Mix Enhancements or Substitutions

List of Additional Species to Add Diversity or for Substitutions of seeds or plugs.

Pollinator Plot Southwest 38-541

Updated 11-10-2020

Grasses:

Scientific Name	Common Name
<i>Bouteloua hirsuta</i>	Hairy Grama
<i>Bromus ciliatus</i>	Fringed Brome
<i>Elymus riparius</i>	Riverbank Wild Rye
<i>Elymus villosus</i>	Downy Wild Rye
<i>Hesperostipa spartea</i>	Porcupine Grass
<i>Koeleria macrantha</i>	Junegrass
<i>Panicum virgatum</i>	Switchgrass
<i>Sorghastrum nutans</i>	Indian Grass

Forbs & Shrubs:

Scientific Name	Common Name
<i>Agastache scrophulariaefolia</i>	Purple Giant Hyssop
<i>Allium canadense</i>	Wild Garlic
<i>Asclepias verticillata</i>	Whorled Milkweed
<i>Boltonia asteroides</i>	False Aster
<i>Chamerion angustifolium</i>	Fireweed
<i>Chelone glabra</i>	White Turtlehead
<i>Doellingeria umbellata</i>	Flat-topped Aster
<i>Drymocallis arguta</i>	Tall Cinquefoil
<i>Galium boreale</i>	Northern Bedstraw
<i>Heterotheca villosa</i>	Hairy Golden Aster
<i>Heuchera richardsonii</i>	Alumroot
<i>Liatris punctata</i>	Dotted Blazing Star
<i>Lobelia spicata</i>	<i>Rough-spiked Lobelia</i>
<i>Lysimachia ciliata</i>	Fringed Loosestrife

<i>Lysimachia quadriflora</i>	Prairie loosestrife
<i>Mimulus ringens</i>	Blue Monkey Flower
<i>Monarda punctata</i>	Horsemint
<i>Pediomelum argophyllum</i>	Silverleaf Scurfpea
<i>Pediomelum esculentum</i>	Prairie Turnip
<i>Penstemon gracilis</i>	Slender Beard Tongue
<i>Physostegia virginiana</i>	Obedient Plant
<i>Polemonium reptans</i>	Spreading Jacob's Ladder
<i>Ranunculus fasciculatis</i>	Early Buttercup
Prairie coneflower	<i>Ratibida columnifera</i>
<i>Rosa arkansana</i>	Prairie rose
<i>Sisyrinchium montanum</i>	Mountain Blue-eyed Grass
<i>Solidago nemoralis</i>	Gray Goldenrod
<i>Solidago ptarmicoides</i>	Upland White Aster
<i>Solidago riddellii</i>	Riddell's Goldenrod
<i>Symphyotrichum oolentangiense</i>	Skyblue Aster
<i>Teucrium canadense</i>	Germander
<i>Thalictrum dasycarpum</i>	Tall Meadow-Rue
<i>Tradescantia ohioensis</i>	Ohio Spiderwort
<i>Vernonia fasciculata</i>	Bunched Ironweed
<i>Zizia aptera</i>	Heart-leaved Alexanders

Legumes:

Scientific Name	Common Name
<i>Astragalus crassicaarpus</i>	Ground Plum
<i>Chamerion angustifolium</i>	Fireweed
<i>Glycyrrhiza lepidota</i>	Wild Licorice
<i>Vicia americana</i>	American Vetch

Bareroot plants or plugs to supplement your planting

Scientific Name	Common Name	Bloom Time	Sun/Shade	Range
<i>Anemone patens</i>	pasqueflower	e	Sun, Part Shade	NW,SW,SE
<i>Antennaria neglecta</i>	pussytoes	e/m	Sun, Part Shade	NW,SW,SE,NE
<i>Cirsium flodmanii</i>	Flodman's Thistle	m/l	Sun	NW, SW, SE, NE
<i>Galium boreale</i>	Northern bedstraw	m/l	Sun, Part Shade	NW,SW,SE,NE

<i>Gentiana puberulenta</i>	Downy gentian	l	Sun	NW, SW, SE
<i>Geranium maculatum</i>	wild geranium	e/m	Sun, Part Shade, Shade	NW,SW,SE,NE
<i>Geum triflorum</i>	Prairie smoke	e/m	Sun	NW,SW,SE, NE
<i>Heuchera richardsonii</i>	Alumroot	e/m	Sun, Part Shade	NW,SW,SE,NE
<i>Lithospermum canescens</i>	Hoary puccoon	e/m	Sun, Part Shade	NW,SW,SE,NE
<i>Pediomelum esculentum</i>	Prairie turnip	e/m	Sun	NW,SW, SE
<i>Rosa arkansana</i>	Prairie rose	m	Sun, Part Shade	NW,SW,SE,NE
<i>Sisyrinchium campestre</i>	Blue-eyed grass	e/m	Sun	NW,SW,SE,NE
<i>Viola palmata var. pedatifida</i>	Prairie violet	e/m	Sun, Part Shade	NW,SW,SE
<i>Viola spp.</i>	Violets	e/m	Part Shade	NW,SW,SE,NE

Bloom Time:

early
Mid
Late

Pollinator Plot Southwest Seed 38-541 Mix Guidance

Seed mix name: Pollinator Plot Southwest 38-541

Geographic area: Minnesota, Southeast

Year of development: 2016

Year/s of update: 2022

Status (*Standard or Pilot mix*): Standard

Primary and Secondary Functions:

Primary – Terrestrial habitat for pollinators and other invertebrates and improve habitat for beneficial soil organisms

Secondary – Carbon sequestration, wildlife habitat, emission reductions,

Similar State Mixes: Pollinator Plot Southeast, Pollinator Plot Northwest, Pollinator Plot Northeast

Compatible NRCS Practice Standards: 327 Conservation Cover (changing to 420 Wildlife Habitat Planting) - Pollinator Species

Compatible Minnesota CRP Practices: CP42

Suitable Site Conditions

Areas with mesic soils and full sun for at least 70% of the day where land is being converted from other uses such as lawn, or agricultural fields like row crops or pasture.



Seed Mix Highlights

This mix has been designed to support specialist bees, many Lepidoptera species and includes a wide range of plant families to maximize insect use, bloom periods and the long-term resiliency of the mix.

How to Customize the Mix for Unique Site Conditions and Goals

Site conditions such as sunlight, soils, hydrology and existing vegetation along with functional goals for the project such as pollinator habitat, carbon sequestration, and benefit to grassland bird species can all inform species selection and the modification of seed mixes. See the Additional Species List, which can be used to amplify diversity. Use the [BWSR Seed Mix Substitution Table](#) when species are not available from vendors, or the landowner has alternative goals for the site.

Site Preparation

The primary goal of site preparation is to control weed species and provide ideal growing conditions for native seeds or plants to be installed. Preparation methods vary depending on the current site conditions. For example, sites with introduced perennial grasses require different suppression techniques than sites that have been farmed with row crops. Specific conditions should be considered when selecting site preparation techniques: unique biology of current (unwanted) vegetation, potential desirable and undesirable species in the seed bank, previous pesticide use on site, and potential for soil erosion. Annual cover crops or row crops can be used in preparation for a native planting, as long as they are not grown with persistent pesticides such as neonicotinoids or other long-lasting chemicals that may negatively impact pollinators or desirable vegetation establishment. For example, following soybean harvest, an un-tilled soybean field provides a good seedbed with potentially little plant residue for planting native prairie mixes. For fields where persistent chemicals have been used (including neonicotinoids or long-lived herbicides) it is recommended to plant a temporary cover crop for one or two seasons to allow the chemicals to break down in the soil. For more details on organic or pesticide-free site preparation techniques, see the Xerces Society guide, [Organic Site Preparation for Wildflower Establishment](#).

Temporary Cover Crops

Short-lived temporary cover crops can help stabilize project sites in preparation for planting native seed mixes. Cover crops such as oats (the most commonly used species) should be mowed to 10-12 inches before seeds mature (or harvested upon maturity) to prevent re-seeding. Other cover crops typically used in agricultural fields, such as buckwheat, winter wheat, sorghum sudangrass, and radishes, can help stabilize soil, build soil health, or provide weed competition as part of restoration projects. Residues from temporary covers should be minimized prior to seeding to increase seed to soil contact.

Seedbed preparation

Methods that are used to prepare a seedbed can vary depending on the type of seeding equipment to be used. If a traditional native seed drill will be used, a smooth, firm seedbed is required. Soybean fields usually are sufficiently prepared for a native seed drill. Corn fields should be raked and bailed if possible or light tillage should be used to remove leftover corn residue that would prevent sufficient seed to soil contact. Sites that were recently tilled will require additional soil treatment such as field cultivation, harrowing and rolling to prepare a firm seedbed and prevent seed from being buried too deep. Native

seeds will have difficulty germinating if they are buried more than ¼" deep. Broadcast seeding can be conducted on soybean or corn fields; fields that have been disked, should be cultipacked or allowed to settle before seeding. Some practitioners have found that broadcast seeding on a smooth surface (not tilled or disked) leads to the establishment of higher diversity. It is important that the soil surface is not too hard packed, so cultipacking or light harrowing of crop fields before broadcast seeding may be needed. Seed can be lost on smooth surfaces, so it is recommended to seed into temporary cover crops or to roll sites after broadcast seeding. BWSR has seedbed preparation guidance based on current site conditions on our website: <https://bwsr.state.mn.us/habitat-establishment-management-resources>

Seeding Methods

A variety of seeding equipment is used for upland pollinator plantings including broadcast seeders, no-till native seed drills, Brillion seeders and Trillion seeders. Specialized native seed drills can handle a wide variety of seed (fluffy, smooth, large and small) and low seeding rates. Conventional grain drills are not capable of handling diverse seed sizes and will not provide satisfactory results. Broadcast seeding is common for planting pollinator mixes. Broadcast seeding equipment should be used that is designed to spread mixes with different sized seeds (e.g. Vicon Seeders). NRCS recommends broadcast seeding at a rate of 1.5 times the normal seeding rate and cultipacking after seeding (based on Agronomy Tech Note 31).

Seeding Dates

Forb-dominated pollinator seed mixes can be installed in the spring or fall but fall dormant plantings allow seeds to naturally stratify and settle into the soil through periods of freezing and thawing over the winter. Spring seedings should be done on or around May 1-July 1 when soil temperatures are at least 60 degrees Fahrenheit or higher. Fall seeding should occur when soil temperatures fall below 50 degrees Fahrenheit for a consistent period of time (usually around October 15 in the northern half of the state and November 1 in the southern half of the state). Fall dormant seedings can help reduce weed pressure during the first year of growth because native cool-season grasses and forbs germinate earlier and start competing with weed species right away. Frost seedings are also an option if the snow cover is shallow, ice-free, and winds are calm. For a frost seeding, seeding rates may need to be increased by 25 -50 percent due to lower germination rates and loss of seed that is consumed by wildlife over the winter months.

Management Methods

Establishment Mowing – Mowing can be an important step in the establishment of pollinator plantings that have high pressure from annual weeds. Expect to mow at least twice during the first season and once during the second season with a flail mower or stalk chopper (to prevent smothering plants) to decrease competition and to provide sufficient sunlight for seedlings. Haying is another method to remove mowed vegetation that prevents smothering of the native seedlings. Mowing should be conducted before weeds mature and seed out. It is important that mowed vegetation does not smother the planting; therefore, very productive sites may need to be mowed more often in the first year to reduce the mulching effect. Vegetation should be mowed to between five and eight inches before seed is allowed to set (usually as weeds reach 12-24 inches). Mowing height should be raised as native plants establish. Mowing too short can be detrimental to the outcomes of a successful planting. Some

grassland managers see success without mowing but the need will vary depending on site conditions (such as soil productivity) and weed pressure.

Spot Mowing – As the native plant community is getting established, it may be beneficial to spot mow or weed trim areas with invasive or noxious plants. Spot-mowing should be done at a raised height between 4-6 inches in order to target the invasive plants and to avoid damaging the nearby native species. Spot mowing for control of invasive or noxious weeds can be done every year to increase the diversity and functionality of the planting. A list of noxious/invasive weed species that should be eradicated can be viewed at the [Minnesota Department of Agriculture's website](#). Mowing is a good way to control some species such as thistles but other methods are needed to control species such as Poison Hemlock, Common Tansy, Leafy Spurge, Spotted Knapweed, Wild Carrot and Wild Parsnip. If there are large areas of these species, it would be helpful to minimize the disturbance of site preparation. There are helpful guidelines in the manual [Restoration-Guide-Invasive-Perennial-to-Conservation-Prairie.pdf \(nature.org\)](#).

Spot Management of Weeds – Some persistent perennial weeds may require digging, pulling, girdling, smothering or spot treatment with herbicides for sufficient control. Some persistent perennial plants include: reed canary grass, smooth brome, quack grass, purple loosestrife, Canada thistle, Kentucky bluegrass, crown vetch, birds-foot trefoil, and woody species, such as box elder, common buckthorn, Siberian elm, and Tartarian honeysuckle. Control methods should be conducted carefully during the early establishment phase, to avoid adverse impacts to native plant seedlings. Herbicides should only be used on persistent perennial weeds; most other weeds will be excluded over time as the native planting matures. Herbicide use should be species specific, sprayed in a discriminating and targeted way (minimizing non-target organisms), and applied according to rates specified on the label. All pesticides should be selected and applied in a manner that minimizes risks to human health, beneficial and nontarget organisms, and the environment. ---

Prescribed Burning – Prescribed burning is beneficial to remove thatch, control invading woody and invasive plants in prairies, fertilize the soil with ashes, stimulate seed germination and new plant growth, and increase diversity in plantings. Burning is typically initiated after the third or fourth years of establishment, as native plants become the dominant cover. Uplands benefit from burning every three to five years. The timing of a burn can also help with management goals. Late spring burns are used to set back cool-season non-native species such as smooth brome and reed canary grass. Fall burns can release spring-blooming species for pollinators. Fall and spring burns should be alternated periodically to simulate natural variation. Burn plans are needed to define the details of how the burn will be conducted, who will be involved and for contingency planning. In many cases, permits are also required. It is recommended to only burn one-third or less of a project site at a time, especially if there is no adjacent refuge or conservation area nearby for wildlife to escape the fire. Partial burns and burns that are patchy may also benefit pollinator populations if timed correctly (when pollinators are not actively foraging or pollinators have pupated and are mobile).

What to Expect in Year 1

During the first year of establishment, many native grasses, sedges and flowers will remain about one to three inches tall. Mowing will help to keep agricultural weeds (foxtail, barnyard grass, ragweed etc.)

managed so the native plant seedlings receive sufficient water and sunlight. The planting may have a somewhat weedy appearance in the first year (see establishment mowing paragraph above).

(IMAGE)

What to Expect in Year 2

During year two, some of the native grasses, sedges and flowers may reach their mature height and some of them may even flower. Depending on when the seeding was conducted (spring or dormant), there might also be many first-year native seedlings germinating alongside native plants that established the year prior. Mowing may still play a key role in managing weeds and allowing seedlings to grow.

(IMAGE)

What to Expect in Year 3 and Beyond

By the end of year three many of the native plants will be mature and should start flowering. There may be some species that are slow to establish and may not show up for several years.

Problem Solving

Poor Establishment After Year 1 – It is often difficult to determine if a seeding is successful during the first year, as establishment may vary depending on weather conditions and some species may be slow to establish. It is typically best to wait until the second year to conduct any corrective actions. Look for species such as Black-Eyed Susan seedlings in year 1 for confirmation the seeding was a success.

Poor Establishment After Year 2 – If native plant seedlings are not establishing about every one to two feet it may be necessary to interseed some species into the planting. Monitor the site during the growing season to determine which species are present, and which species may need to be supplemented. Interseeding should be conducted after the second growing season.

High Annual and Biennial Weed Competition – Typically, annual and biennial weed competition is not a big problem in prairie plantings as they are short lived and frequent mowing in the first year of establishment reduces their contributions to the seed bank. In addition, perennial native plants tend to outcompete annual and biennial weeds as the planting matures.

High Perennial Weed Competition – Dense establishment of perennial species can be a problem as it can prevent the establishment of native species. Prescribed burning, prescribed grazing, and/or spot herbicide application may be needed to manage perennial weeds.

Low Forb Diversity After Year 3 – If grasses and sedges are establishing successfully but there is a lack of forbs it is recommended to interseed additional forbs in late fall or after a prescribed fire in spring or fall. See the [Xerces Society guide](#) for additional information and guidance about interseeding wildflowers.